

## Bibliography

- [1] ALTSHULER I., DEMIRI B., XU S., CONSTANTIN A., YAN N.D., CRISTESCU M.E. An integrated multi-disciplinary approach for studying multiple stressors in freshwater ecosystems: Daphnia as a model organism. *Integr. Comp. Biol.* 2011, **51** pp. 623–633
- [2] MALDONADO-ALAMEDA A., GIRO-PALOMA J., RODRÍGUEZ-ROMERO A., SERRET J., MENARGUES A. Andrés A., Chimenos, JM. Environmental potential assessment of MSWI bottom ash-based alkali-activated binders. *J. Hazard. Mater.* 2021, **416** p. 125828
- [3] RODRIGUES J.D., SILVESTRE J.D., VIEGAS C.A., AHMED J.J., KURDA R., DE BRITO J. Evaluation of the Ecotoxicological Potential of Fly Ash and Recycled Concrete Aggregates Use in Concrete Appl. *Sci.* 2020, **10** p. 351
- [4] SCHMITTGEN T.D., LIVAK K.J. Analysing real-time PCR data by the comparative CT method. *Nat. Protoc.* 2008, **3** pp. 1101–1108
- [5] EN ISO 6341, *Water quality - Determination of the inhibition of the mobility of Daphnia magna Straus (Cladocera, Crustacea) - Acute toxicity test (ISO 6341)*